

ENT. To-morrow, Roey, by steam to Edinburgh ; but let us stay here as long as we can, and now—

Hie away! to the woods, hie away!
 Aurora with crimson has tinted the sky,
 And the blithe lark, the herald of day,
 Is pouring his music around from on high.
 Hie away! to the woods, hie away!

Hie away! to the woods, hie away!
 Field, flower, and forest, all glitter with dew,
 And there droops, on its elegant spray,
 The harebell, arrayed in its beautiful blue.
 Hie away! to the woods, hie away!

It is not the first time that we have made a door of this window.

ERRO. And the other doors will be fastened now. (*He takes up the nets.*)

Et jam prima novo spargebat lumine terras
 Tithoni croceum linquens Aurora cubile:
 Jam sole infuso, jam rebus luce retutis
 Turnus in arma viros, armis circumdatus ipse
 Suscitatur.

(*Exeunt through the open window.*)

ART. XXXIII.—*Monographia Chalciditum.* By FRANCIS WALKER.

(*Continued from p. 309.*)

“ ——— the green myriads in the peopled grass.”

GENUS VIII.—PLATYTERMA.

Sp. 7. Plat. incultum. Mas. P. prasini statura, antennis fuscis, alis subhyalinis.

Viride: oculi ocellique rufo-fusci: antennæ pallidè fuscae, corporis dimidio longiores, subtus fulvæ; articulus 1^{us}. lætè flavus; clava obscurè fusca: abdomen thorace paullò longius; discus obscurè cupreus: sexualia fusca: pedes lætè flavi; coxæ virides; tarsi apice fusci; protarsi fulvi: alæ subhyalinæ; squamulæ et nervi flava; stigma pallidè fuscum, minutum. (Corp. long. $\frac{5}{4}$ lin.; alar. $1\frac{1}{4}$ lin.)

September; near the Land's End, Cornwall.

Sp. 8. Plat. comptum. Fem. P. terminali *simile*, caput et abdomen latiora, antennæ obscuriores.

Lætè æneo-viride : caput thorace latius : oculi ocellique rufo-fusci : antennæ fuscæ, corporis dimidii longitudine ; articulus 1^{us}. basi flavus : metathorax viridis : abdomen viride, elongato-ovatum, thorace paullò longius et latius, subtus non angulatum ; discus cupreo-æneus : pedes pallidè flavi ; coxæ virides ; tarsi apice fusci ; protarsi flavi : alæ hyalinæ ; squamulæ et nervi pallidè flava ; stigma minutum. (Corp. long. 1 lin. ; alar. 1½ lin.)

July ; on grass in fields ; near London.

Sp. 9. Plat. femorale. Mas et Fem. *Viride*, antennis *fulvis apice fuscis* (mas) *aut fuscis* (fem.), *pedibus flavis*, fem. *femoribus viridibus*, *alis hyalinis*.

Mas.—Lætè viride : caput thoracis latitudine : oculi ocellique rufo-fusci : antennæ fulvæ, corporis dimidii longitudine ; articulus 1^{us}. flavus ; 2^{us}. supra et clava fusca : abdomen æneo-viride, nitens : sexualia fulva : pedes lætè flavi ; coxæ virides ; tarsi apice fusci ; protarsi fulvi : alæ hyalinæ, paullò flavescentes ; squamulæ et nervi flava ; stigma minutum.

Fem.—Viride : antennæ fuscæ ; articulus 1^{us}. fulvus ; 2^{us}. viridis, apice fuscus : abdomen lætè viride, thorace vix longius, subtus non angulatum ; segmenta apice cuprea : femora viridia, apice flava ; meso- et metatarsi apice nigro-fusci : alæ fulvescentes ; squamulæ et nervi fulva ; stigma pallidè fuscum, minutum. (Corp. long. ½—1 lin. ; alar. ½—1½ lin.)

Var. β.—*Mas*, antennæ articulis 11^o. basi et 13^o. apice fulvis.

Var. γ.—*Mas*, caput cyaneo-viride.

Var. δ.—*Mas*, antennæ clava fulva.

Var. ε.—*Mas*, abdomen viride.

Var. ζ.—*Mas*, abdomen viride ; discus cupreo-viridis.

Var. η.—*Mas*, cyaneo-viride : abdominis discus æneo-viridis.

Var. θ.—*Mas*, abdominis discus cupreus.

Var. ι.—*Mas*, caput et thorax cyaneo-viridia, illum inter oculos et hujus scutum posticè æneo-viridia ; abdomen viride ; discus cupreus.

Var. κ.—*Fem.* antennæ articulo 1^o. supra fusco.

Var. λ.—*Fem.* antennæ articulo 2^o. viridi-fusco.

Var. μ.—*Fem.* antennæ articulo 1^o. supra viridi-fusco.

Var. ν.—*Fem.* abdominis discus cupreo-æneus.

Var. ξ.—*Fem.* abdominis segmenta apice cyaneo-cuprea.

Var. ο.—*Fem.* femora et tibiæ fulva.,

Var. π.—*Fem.* abdomen cupreo-æneum, basi apice que viride; segmenta nonnulla basi cyanea: alæ vix fulvescentes.

Var. ρ.—*Fem.* alæ omninò perlucidæ.

Var. σ.—*Fem.* thoracis dorsum æneo-viride: stigma fuscum.

Var. τ.—*Fem.* thoracis latera, abdomen subtus et femora æneo-viridia.

April to September; on grass in fields; near London. June; Windsor. June and September; Isle of Wight. September; Westmoreland and Cumberland; Penzance, Cornwall. New Lanark, Scotland.

Sp. 10. Plat. decorum. Mas et Fem. *Viride, abdomine purpureo, antennis fulvis (mas) aut fuscis (fem.), pedibus flavis, alis griseo-hyalinis.*

Mas.—Læte aureo-viride: caput thorace vix latius: oculi ocellique rufo-fusci: antennæ fulvæ, corporis dimidio paullò breviores; articulus 1^{us}. lætè flavus, apice fulvus; 2^{us}. basi pallidè fuscus; clava obscurè fusca, apice pallidior, articulo 10^o. paullò latior: caput anticè mesothoracisque scutellum apice viridia: abdomen thorace paullò longius, basi cupreo-viride; discus splendide cupreo-purpureus: sexualia pallidè fusca: pedes lætè flavi; coxæ virides; tarsi apice fusci; protibiæ et protarsi fulva; alæ griseo-hyalinæ; squamulæ flavæ; nervi fusci; stigma minutum.

Fem.—Obscurè viride: antennæ fulvo-fuscæ, corporis dimidio multò breviores; articulus 1^{us}. fulvus; 2^{us}. fuscus: abdomen obscurè purpureum, thorace multò longius et paullò angustius, basi lætè cupreo-viride, subtus æneo-viride angulatum, apice concolor attenuatum acuminatum pubescens. (Corp. long. $1\frac{1}{4}$ — $1\frac{1}{3}$ lin.; alar. $1\frac{1}{2}$ — $1\frac{3}{4}$ lin.)

July and August; on oak trees; near London.

Sp. 11. Plat. remotum. Fem. *Præcedentibus latius et brevius.*

Lætè viride: caput thoracis latitudine: oculi ocellique rufo-fusci: antennæ fulvæ, corporis dimidii longitudine; articulus 1^{us}. pallidior; clava obscurior: parapsidum suturæ æneo-virides: abdomen obscurè cupreo-æneum, thoracis longitudine et latitudine, basi lætè viride, subtus angulatum, apice acuminatum; segmenta basi viridi-ænea: pedes lætè flavi; coxæ virides; femora basi, meso- et metatibiæ medio pallidè fusca: tarsi apice fusci: protibiæ et protarsi fulva: alæ hyalinæ, minimè sub costam flaves-

centes; squamulæ et nervi flava; stigma minutum. (Corp. long. $\frac{4}{5}$ lin.; alar. $1\frac{1}{5}$ lin.)

July; on grass in fields; near London.

GENUS IX.—AMBLYMERUS.

Sp. 4. Amb. ruralis. Fem. A. valido *paullò minor et angustior; alæ obscuriores.*

Æneo-viridis: caput viride, thorace paullò latius: oculi ocellique rufo-fusci: antennæ fuscæ; articulus 1^{us}. flavus: abdomen obscurè cupreum, thorace paullò longius, subtus angulatum; segmentorum margines basi et utrinque lætè virides: oviductus pallidè rufus, abdominis apicem non transiens: pedes pallidè fulvi; coxæ æneo-virides; genua, tibiæ apice tarsique pallidè flava, hi apice fusci; protarsi flavi: alæ subfulvæ; squamulæ et nervi fulva; stigma minutum. (Corp. long. $\frac{5}{4}$ —1 lin.; alar. $1-1\frac{1}{4}$ lin.)

Var. β.—Antennæ articulo 1^o. fusco, basi subtusque flavo.

Var. γ.—Abdomen ferè omne nigro-cupreum; segmentum 1^{um}. basi lætè viride.

August; on grass beneath trees; near London. September; Isle of Wight; New Lanark, Scotland.

Sp. 5. Amb. campestris. Fem. *Præcedenti similis, paullò brevior; clava angustior et acutior.*

Æneo-viridis, vix nitens: caput viride, thorace paullò latius: oculi ocellique rufo-fusci: antennæ fuscæ; articulus 1^{us}. flavo-fulvus, basi flavus: abdomen nigro-cupreum, thorace vix longius, subtus non angulatum; segmenta basi lætè viridia: pedes fulvi; coxæ æneo-virides; protibiæ pallidè fulvæ, subtus flavæ; tarsi apice fulvi; meso- et metatarsi straminei: alæ subfulvæ; squamulæ et nervi fulva; stigma parvum. (Corp. long. $\frac{1}{2}$ lin.; alar. $\frac{5}{4}$ lin.)

Var. β.—Protibiæ omninò flavæ: meso- et metapedum genua tibiæque apice straminea.

August; on grass in fields; near London.

Sp. 6. Amb. latus. Fem. *Viridis, A. dubii statura, clava angustiore, antennis fuscis, pedibus flavis, alis subhyalinis.*

Viridis: caput thoracis latitudine: oculi ocellique rufi: antennæ obscurè fuscæ; clava pallidior; articulus 1^{us}. fulvus: abdomen

thorace paullò longius, subtus non angulatum; discus nigro-æneus; segmenta 2^o. ad 4^{um}. cyaneo cingulata: pedes lætè flavi: coxæ virides; meso- et metatarsi straminei, apice fusci; protarsi apice fulvi: alæ subhyalinæ, vix flavescentes; squamulæ fuscae; nervi flavi; stigma parvum. (Corp. long. 1 lin.; alar. 1½ lin.)

September; Isle of Wight.

Sp. 7. Amb. truncatellus. Fem. *Æneo-viridis, præcedentibus multò brevior, antennis obscurè fuscis, pedibus flavis, alis hyalinis.*

Æneo-viridis: caput thorace paullò latius: antennæ obscurè fuscae, corporis dimidio vix longiores; clava pallidior; articulus 1^{us}. basi flavus: oculi ocellique fusco-rufi: abdomen viride, basi nitentius et cupreo variegatum, thorace paullò angustius vix longius, subtus angulatum; discus æneus: pedes flavi; coxæ æneo-virides; metafemora extus et mesotibiæ intus fulva; meso- et metatarsi straminei, apice fusci; protarsi apice fulvi: alæ hyalinæ; squamulæ et nervi flava; stigma minutum. (Corp. long. ½ lin.; alar. ¾ lin.)

July; on grass in fields; near London.

Sp. 8. Amb. fulvipennis. Fem. *Cupreo-æneus, ferè A. validi statura, antennis fuscis, pedibus fulvis, alis fulvis.*

Cupreo-æneus: caput viridi-æneum, thorace paullò latius: oculi ocellique rufo-fusci: antennæ obscurè fuscae, corporis dimidio vix breviores; articulus 1^{us}. fulvus: abdomen cupreum, breve, latum, thoracis longitudine, subtus angulatum; discus obscurior: pedes fulvi; coxæ cupreo-æneæ; meso- et metatarsi flavi, apice fusci: alæ fulvæ; squamulæ et nervi fusca; stigma minutum. (Corp. long. 1 lin.; alar. 1½ lin.)

Var. β.—Abdomen basi viridi-æneum: tibiæ apice basique flavæ: squamulæ et nervi pallidè fusca.

June; on grass beneath trees; near London; New Forest, Hampshire.

Sp. 9. Amb. modestus. Fem. *Viridis, præcedentis ferè statura sed minor, antennis nigro-fuscis, abdomine plerumque æneo, pedibus fulvis, alis fuscis.*

Obscurè viridis: caput thorace latius: oculi ocellique rufo-fusci: antennæ nigro-fuscae, crassæ, corporis dimidio breviores; articulus

1^{us}. fulvus: abdomen viridi-æneum, thorace non longius sed multò latius, subtus non angulatum, basi nitentius: oviductus rufus: pedes fulvi; coxæ virides; femora fusca, apice fulva; meso- et metatarsi flavi, apice fuscì: alæ fuscae; squamulae et nervi fulvo-fusca; stigma parvum. (Corp. long. $\frac{1}{2}$ — $\frac{5}{4}$ lin.; alar. $\frac{3}{4}$ —1 lin.)

Var. β.—Antennæ fuscae; articulus 1^{us}. fulvus, apice fuscus.

Var. γ.—Antennæ subnigræ; articulus 1^{us}. obscurè fuscus, basi fulvus.

Var. δ.—Antennæ articulo 1^o. omninò fusco.

Var. ε.—Antennæ articulo 1^o. apice fusco: abdomen nigro-viride, basi æneo-viride nitentius.

Var. ζ.—Antennæ articulo 1^o. fulvo-fusco: abdomen viride; discus nigro-viridis: pedes flavi; femora fulva; meso- et metatarsi straminei, apice fuscì: alæ subfuscae.

Var. η.—Antennæ articulo 1^o. fusco, basi fulvo: abdomen obscurè æneo-viride; discus cyaneo-cupreus: alæ subfuscae.

Var. θ.—Abdomen cupreo-æneum, basi viride nitentius, apice æneo-viride.

Var. ι.—Abdomen nigro-æneum, basi æneo-viride nitentius: femora omninò fulva.

Var. κ.—Abdomen basi æneo-viride: femora fulva; protibiæ et protarsi flava.

Var. λ.—Cyaneo-viridis: abdomen obscurè viridi-cyaneum, basi æneo-viride.

Var. μ.—*Var. λ* similis: femora fulva: alæ subfuscae.

Var. ν.—Antennæ articulo 1^o. flavo: protibiæ et protarsi flava: abdomen æneo-viride, basi viride: femora fulva.

Var. ξ.—Abdomen nigro-æneum, basi apiceque viride: femora fulva; protibiæ et protarsi flava: alæ fuscae; discus obscurior.

Var. ο.—Obscurè viridi-æneus: abdomen nigro-æneum, basi apiceque æneo-viride: femora fulva.

Var. π.—Abdominis segmenta basi nigro-cyanea: femora fulva; alæ subfuscae.

Var. ρ.—Antennæ fuscae; articulus 1^{us}. apice fuscus: abdomen viride; discus cyaneo-cupreus: pedes flavi; meso- et metatarsi straminei, apice fuscì: alæ subhyalinae.

Var. σ.—Æneo-viridis: antennæ nigræ; articulus 1^{us}. nigro-fuscus, basi fulvus: abdominis segmenta apice nigro-cyanea: pedes fuscì; tarsi flavi, apice fuscì; genua et protarsi fulva: alæ subfuscae.

August; on grass in fields; near London. September; Isle of Wight; Westmoreland and Cumberland; Land's End, Cornwall; New Lanark, Scotland.

Sp. 10. Amb. fuscipes. Fem. *Viridi-æneus*, A. amoeni statura, abdomine cupreo-æneo, antennis pedibusque fuscis, alis fulvescentibus.

Viridi-æneus: caput viride, thorace latius: oculi ocellique rufo-fusci: antennæ fuscae, sat crassæ, corporis dimidio vix breviores; articulus 1^{us}. obscurè fuscus, basi pallidior: abdomen purpureo-æneum, thorace paullò longius vix latius, apice basique æneo-viride, subtus angulatum et apice minimè elevatum; segmenta basi viridia: pedes obscurè fusci; coxæ virides; mesofemora subtus apice spina armata; genua fulva; protarsi fusci; meso- et metatarsi pallidè flavi, apice fusci: alæ subhyalinæ, amplæ, sub nervum fulvæ; squamulæ fuscae; nervi fulvi; stigma parvum. (Corp. long. 1 lin.; alar 1 $\frac{3}{4}$ lin.)

Var. β .—Caput supra æneum: thorax et abdomen cupreo-ænea, hujus discus cyaneo-purpureus: protarsi fulvi, apice fusci.

June; on grass in woods; near London.

Sp. 11. Amb. humilis. Fem. *Viridi-æneus*, præcedenti similis, antennis pedibusque fuscis, femoribus viridibus, abdomine cupreo, alis subflavescentibus.

Æneus: caput æneo-viride: oculi ocellique rufo-fusci: antennæ fuscae; articulus 1^{us}. obscurior, basi fulvo-fuscus: abdomen obscurè cupreum, basi et subtus æneo-viride nitentius; segmenta basi viridia: pedes fusci; coxæ et femora viridia; protibiæ pallidè fuscae; genua et protarsi fulva; meso- et metatarsi flavi, apice fusci: alæ hyalinæ, sub costam flavescentes; squamulæ et nervi fulva; stigma obscurius, parvum. Corp. long. $\frac{4}{5}$ lin.; alar. 1 $\frac{1}{2}$ lin.)

June; New Forest, Hampshire.

Sp. 12. Amb. albitarsus. Fem. *Viridi-cyaneus*, A. fuscipedis statura, abdomine purpureo, antennis pedibusque fuscis, alis hyalinis.

Viridi-cyaneus: caput thorace latius: oculi ocellique rufo-fusci: antennæ fuscae, graciles, corporis dimidii longitudine: abdomen purpureum, thorace longius et angustius, apice obscurè basi lætè viride, subtus non angulatum: metathorax lætè viridis: pedes nigro-fusci; coxæ virides; genua, tibiæ apice tarsique alba, hi apice nigro-fusci; propedum genua flava, tibiæ tarsique pallidè fusca: alæ hyalinæ, latæ; squamulæ fuscae; nervi fulvi; stigma parvum. (Corp. long. 1 lin.; alar. 1 $\frac{1}{5}$ lin.)

August; on grass beneath trees; near London.

Sp. 13. Amb. nitescens. Fem. *Viridi-æneus*, *A. amœni statura, abdomine purpureo-æneo, antennis pedibusque nigro-fuscis, alis subfuscis.*

Æneus: caput thorace latius, anticè viride: oculi ocellique rufo-fusci: antennæ nigro-fuscae, corporis dimidio paullò longiores; articulus 1^{us}. fuscus, basi fulvus: metathorax æneo-viridis: abdomen viride, thorace longius, subtus non angulatum; discus purpureo-æneus; segmenta basi viridia: coxæ virides; trochanteres fulvi; femora nigro-viridia; tibiæ nigrae, apice basique flavæ; protibiæ et protarsi fusca, illæ extus nigrae; meso- et metatarsi flavi, apice fusci; alæ subfuscae, prope costam obscuriores; squamulae et nervi fusca; stigma mediocrem, nigrum. (Corp. long. $1\frac{1}{2}$ lin.; alar. $2\frac{1}{2}$ lin.)

August; on grass under trees; near London.

Sp. 14. Amb. pusillus. Fem. *Æneo-viridis, præcedentibus angustior, A. dubio similis at minor, antennis fuscis, pedibus flavis, alis subhyalinis.*

Æneo-viridis: caput thorace paullò latius: oculi ocellique fuscorufi: antennæ fuscae, corporis dimidii longitudine; articulus 1^{us}. basi flavus: scutellum viridi-æneum: abdomen thorace longius, subtus non angulatum; discus cupreo-æneus: pedes flavi; coxæ æneo-virides; tarsi apice fusci: alæ subhyalinae, minimè flavescentes; squamulae fulvae; nervi flavi; stigma minutum. (Corp. long. $\frac{1}{2}$ — $\frac{2}{5}$ lin.; alar. $\frac{3}{4}$ —1 lin.)

Var. β.—Caput viride.

Var. γ.—Viridis: prothorax æneo-viridis: abdominis discus cupreo-æneus.

September; Isle of Wight.

Sp. 15. Amb. tenuicornis. Fem. *A. pusillo similis, antennis gracilioribus, abdomine longiore et angustiore.*

Æneo-viridis: caput anticè viride, thoracis latitudine: oculi ocellique rufo-fusci: antennæ fuscae, corporis dimidii longitudine, hujus generis plerisque graciliores clava angustiore et acutiore; articulus 1^{us}. basi flavus: abdomen viride, thorace longius, subtus paullò carinatum; segmenta 2^o. ad 5^{um}. nisi ad apices nigro-ænea: pedes flavi; coxæ æneo-virides; tarsi apice pallidè fusci: alæ hyalinae; squamulae et nervi flava; stigma minimum, vix conspicuum. (Corp. long. $\frac{2}{3}$ lin.; alar. $\frac{3}{4}$ lin.)

July; on grass in fields; near London.

Sp. 16. Amb. hebes. Fem. *Viridis*, A. modesti *statura*, *præcedentibus plerisque crassior*, *antennis fuscis*, *abdomine æneo*, *pedibus fulvis*, *alis hyalinis*.

Viridis: caput thorace vix latius: oculi ocellique fusci: antennæ obscurè fuscæ, crassæ, corporis dimidii longitudine; articulus 1^{us}. flavus, apice fulvus: scutellum æneo-viride: abdomen thorace vix longius, subtus non angulatum, medio cupreo-æneum parùm nitens, basi æneum nitentius: pedes fulvi; coxæ æneo-virides; tarsi flavi, apice fusci: alæ hyalinæ; squamulæ et nervi fusca; stigma minutum. (Corp. long. $\frac{3}{4}$ lin.; alar. 1 lin.)

August; on grass in fields; near London.

Sp. 17. Amb. tenellus. Fem. A. pusillo *similis sed paullo longior et angustior*.

Viridis: caput thorace vix latius: oculi ocellique fusco-rufi: antennæ fuscæ, corporis dimidio breviores; articulus 1^{us}. basi flavus: abdomen thorace multò longius, subtus non angulatum; discus cupreo-æneus: pedes flavi; coxæ virides; tarsi pallidè flavi, apice fusci: alæ subhyalinæ; squamulæ et nervi fulva; stigma parvum. (Corp. long. $\frac{2}{3}$ lin.; alar. $\frac{3}{4}$ lin.)

September; Isle of Wight.

Sp. 18. Amb. fulvipes. Fem. *Præcedenti similis*, *clava latiore*, *abdomine subtus angulato*.

Viridis: caput thorace paullo latius: oculi ocellique rufi: antennæ pallidè fuscæ, corporis dimidio vix breviores; articulus 1^{us}. basi flavus; clava articulo 10°. multò latior: scutellum æneo-viride: abdomen æneo-viride, thorace paullo longius, subtus angulatum: pedes fulvi; coxæ virides; protibiæ et tarsi omnes flava, hi apice fusci: alæ subfulvæ; squamulæ et nervi fulva; stigma minutum. (Corp. long. $\frac{2}{3}$ lin.; alar. $\frac{3}{4}$ lin.)

September; Isle of Wight.

Sp. 19. Amb. stupidus. Fem. A. lato *similis sed paullo angustior*.

Viridis: caput thoracis latitudine: oculi ocellique rufo-fusci: antennæ fuscæ, corporis dimidio breviores; articulus 1^{us}. basi flavus: abdomen æneo-viride, thorace longius, subtus angulatum: segmenta 2°. ad 5^{um}. apice cupreo-ænea: pedes flavi; coxæ virides; femora viridia, apice basique flava; meso- et metatarsi pallidè

flavi; omnes apice fusci: alæ subhyalinæ, paullò flavescentes; squamulæ et nervi fulva; stigma parvum. (Corp. long. $\frac{2}{3}$ —1 lin.; alar. 1—1 $\frac{1}{2}$ lin.)

Var. β .—Tibiæ fulvæ.

Var. γ .—Thoracis dorsum æneo-viride: protarsi fulvi.

Var. δ .—Antennæ articulo 2^o. viridi-fusco.

Var. ϵ .—Caput, thorax, et femora viridi-ænea.

September; Isle of Wight.

Sp. 20. Amb. nanus. Fem. A truncatelli *ferè statura, paullò angustior.*

Viridis: caput supra cyaneo-viride, thorace paullò latius: oculi ocellique rufo-fusci: antennæ fulvæ, corporis dimidii longitudine; articulus 1^{us}. basi flavus: abdomen viridi-æneum, thorace vix longius, subtus angulatum: pedes flavi; coxæ virides; femora et metatibiæ fusca; mesotibiæ fulvæ; meso- et metatarsi pallidè straminei, apice fusci: alæ hyalinæ; squamulæ et nervi pallidè fulva; stigma minutum. (Corp. long. $\frac{1}{2}$ lin.; alar. $\frac{3}{4}$ lin.)

July; on grass in fields; near London.

Sp. 21. Amb. linearis. Fem. A. tenuicornis *statura, antennis apice crassioribus, clava brevior et obtusior, alis subfulvescentibus.*

Viridi-æneus: caput thoracis latitudine: oculi ocellique rufo-fusci: antennæ fuscae, corporis dimidio breviores; articulus 1^{us}. basi flavus: mesothoracis scutellum et scuti discus cupreo-ænea: abdomen æneum, thoracis longitudinem superans, subtus paullò angulatum, basi cupreo-æneum nitentius; discus nigro-æneus: pedes fulvi; coxæ viridi-æneæ; femora et tibiæ apice, meso- et metatarsi flava, hi apice fusci; protarsi apice obscurè fulvi: alæ subfulvæ, apice et posticè pallidiores; squamulæ et nervi pallidè fulva; stigma minutum. (Corp. long. $\frac{1}{2}$ — $\frac{2}{3}$ lin.; alar. $\frac{2}{3}$ — $\frac{3}{4}$ lin.)

Var. β .—Æneo-viridis: scutellum æneum: abdomen nigro-æneum, basi æneo-viride nitentius; segmenta basi et utrinque viridia: metafemora subfusca.

Var. γ .—Æneo-viridis: caput viride: abdominis discus cupreo-æneus.

September; on grass in fields; near London. Isle of Wight.

Sp. 22. Amb. temperatus. Fem. *Præcedenti similis, antennis paullo brevioribus et crassioribus, abdomine latiore.*

Viridis: caput thoracis latitudine: oculi ocellique rufi: antennæ fuscæ, corporis dimidio breviores; articulus 1^{us}. fulvus, apice supra fuscus: abdomen nigro-æneum, thorace longius, subtus non angulatum, apice basique viride: pedes fulvi; coxæ virides; tarsi apice fusi; genua, meso- et metatarsi flava: alæ subhyalinæ, vix flavescentes; squamulæ et nervi pallidè fulva: stigma minutum. (Corp. long. $\frac{2}{3}$ lin.; alar. $\frac{3}{4}$ lin.)

Var. β .—Æneo-viridis: abdomen nigro-æneum, apice basique æneo-viride.

July; on grass in fields; near London. September; Isle of Wight.

Sp. 23. Amb. iners. Fem. *Præcedenti similis sed latior, antennis brevioribus et crassioribus.*

Viridi-æneus: caput thoracis latitudine: oculi ocellique rufo-fusci: antennæ fuscæ, corporis dimidio breviores; articulus 1^{us}. flavus: abdomen nigro-æneum, thorace latius et paullò longius, basi apiceque viride, subtus non angulatum: pedes fulvi; coxæ æneo-virides; tarsi flavi, apice fusi: alæ fulvæ; squamulæ et nervi fulva; stigma parvum. (Corp. long. $\frac{1}{2}$ lin.; alar. $\frac{2}{3}$ lin.)

September; Isle of Wight.

Sp. 24. Amb. trossulus. Fem. A. modesti *statura, abdomine paullo brevior, antennis fulvis, alis hyalinis.*

Æneo-viridis: caput viride, thorace paullò latius: oculi ocellique rufo-fusci: antennæ fulvæ, crassæ, corporis dimidio longiores; articulus 1^{us}. flavus; 2^{us}. supra basi et clava pallidè fusca: abdomen obscurè cupreum, thorace latius sed non longius, basi lætè viride, subtus non angulatum; segmenta apice viridi-ænea: pedes flavi; coxæ virides; meso- et metatarsi straminei, apice pallidè fusi: alæ hyalinæ, minimè flavescentes; squamulæ pallidè fuscæ; nervi flavi; stigma fulvum, parvum. (Corp. long. $\frac{2}{3}$ lin.; alar. $\frac{3}{4}$ lin.)

August; on grass in fields; near London.

Sp. 25. Amb. stenomerus. Fem. A tenuicorni *similis, paullò brevior et gracilior, alis angustioribus.*

Viridis, angustus, sublinearis: caput thorace paullò latius: oculi ocellique rufofusci: antennæ pallidè fuscæ, graciles, corporis dimidii vix longitudine; articulus 1^{us}. fulvus; clava articulo 10°. multò latior: scutellum et abdomen viridi-ænea; hoc thorace

paullò longius, subtus non angulatum: pedes pallidè fusci; coxæ virides; protibiæ et tarsi omnes flava, hi apice fusci: alæ subfuscæ, angustæ; squamulæ et nervi fulva; stigma minutum. (Corp. long. $\frac{2}{3}$ lin.; alar. $\frac{3}{4}$ lin.)

October; on grass beneath trees; near London.

Sp. 26. Amb. tenebricus. Fem. A. nitescentis *statura, obscurior, stigmatè majore.*

Viridi-æneus, obscurus, crassus: caput æneo-viride, thoracis latitudine: oculi ocellique fusci: antennæ nigræ, crassæ, corporis dimidio paullò breviores; articulus 1^{us}. fuscus, basi fulvus: mesothoracis scutellum cupreo-æneum: abdomen obscurè cupreum, thorace brevius et paullò latius, subtus angulatum, basi cupreo-viride nitentius, apice elevatum: pedes nigro-fusci; coxæ et femora viridi-ænea; trochanteres fusci; genua fulva; tarsi articulo 1^o. fusco, basi pallidè flavo: alæ obscurè fuscæ; squamulæ fulvæ; nervi nigro-fusci, metalis pallidiores; stigma nigrum, magnum. (Corp. long. $1\frac{1}{2}$ lin.; alar. 2 lin.)

October; on grass in fields; near London.

Sp. 27. Amb. mirus. Fem. *Viridiæneus, antennis nigris, abdomine basi pedibusque fulvis, oviductu exerto, alis hyalinis.*

Nigro-viridis, quasi productus, *Callimomi* similis: caput thorace paullò latius: oculi ocellique fusci: trophi ferruginei: antennæ nigræ, crassæ, breves, corporis trientis longitudine; articuli 1^o. ad 5^{um}. fulvi, clava nigro-fusca, ferè rotunda: prothorax lætè viridis, macula supra cupreo-cyanea: thorax subtus viridi-cyaneus: abdomen cupreo-æneum, thorace paullò longius, subtus angulatum, basi fulvum, apice fuscum; latera æneo viridia: oviductus rufus, abdominis trientis longitudine; vaginæ nigro-fuscæ, pilosæ, apice pallidiores: pedes fulvi; coxæ cyaneæ, apice flavæ; femora extus pallidè fusca; trochanteres, genua, meso et metatarsi flava, hi apice fulvi: alæ hyalinæ angustæ; squamulæ et nervi fulva; stigma parvum. (Corp. long. $1\frac{1}{4}$ lin.; alar. $1\frac{3}{4}$ lin.)

August; on grass in fields; near London.

†† *Antennæ articulis 3^o. et 4^o. minimis, 5^o. mediocri.*

§ *Clava brevis, vix acuminata.*

Mari mesotibiæ	{	-intus dilatata	PLATYMESOPUS.
		-extus dilatata	MESOPOLOBUS.
		-simplices	EUTELUS.

GENUS X.—PLATYMESOPUS, *Westwood*.

Mas.—Caput mediocre, thorace paullò latius: antennæ clavatæ; articulus 1^{us}. validus, subfusiformis; 3^{us}. et 4^{us}. minimi; 5^{us}., 6^{us}. et 7^{us}. subæquales; 8^{us}., 9^{us}. et 10^{us}. gradatim latitudine crescentes; clava ovata, articulo 10^o. multò latior: mandibulæ 4-dentatæ, subquadratæ, vix arcuatæ, intùs emarginatæ, subæquales; una dente externo mediocri acuminato, 2^o. obtusiore et multò minore, 3^o. et 4^o. minutis, hoc latiore et obtusiore; altera dentibus brevioribus, 3^o. et 4^o. minimis vix discretis aut conspicuis: maxillæ basi latæ, inde abruptè angustatæ, angulum utrinque extus fingentes, ad apices elongatæ acuminatæ; lacinia in lobum quæque intus producta; palpi 4-articulati, crassi, longitudine mediocri; articuli 1^{us}. et 2^{us}. breves; 3^{us}. et 4^{us}. magni, dilatati, hic ovatus, ille ferè rotundus: labium ^a angustum, sublineare, posticè conicum; ligula ^b parva, anticè quasi fissa; palpi 3-articulati, breves, crassi; articulus 2^{us}. brevissimus; 3^{us}. apice acuminatus: thorax ovatus: prothorax brevissimus: mesothoracis scutum et scutellum maxima; parapsidum suturæ vix conspicuæ; paraptera et epimera benè determinata: metathorax parvus, subtilissimè squameus: abdomen elongato ovatum, thorace angustius sed non longius; segmentum 1^{um}. magnum; sequentia breviora, subæqualia: sexualia exerta: pro- et metapedes simplices; mesopedum femora apices versus spina gracili subtus armata, tibiæ intus valde dilatatæ: alæ mediocres; nervus humeralis ulnari vix longior, setis armatus, ramulum rejiciens nullum; cubitalis radiali brevior, stigmate ramulum brevissimum emittente terminatum.

Fem.—Antennæ articulo 1^o. gracili, 5^o. et sequentibus ad 10^{um}. gradatim latitudine crescentes; clava articulo 10^o. paullò latior; palpi maxillares graciles, filiformes; articuli 1^o. ad 3^{um}. subæquales; 4^{us}. 2ⁱ. et 3ⁱ. longitudine: abdomen thorace paullò latius, subtus non angulatum, apice acuminatum: oviductus non exertus: mesopedum tibiæ simplices.

Sp. 1. *Platy. tibialis. Mas. Viridis, antennis fulvis apice nigris, abdomine cupreo, pedibus flavis, alis hyalinis. Fem. Viridi-æneus, cupreo variegatus, antennis pedibusque fuscis, alis subhyalinis.*

^a The *mentum* of my former descriptions. Vide *Osteology of Insects*, by Mr. Newman, page 71 of this volume.

^b The *labium* of my former descriptions.

Platymesopus tibialis. *Westwood, Lond. and Edinb. Phil. Mag. Third Series. Vol. II. No. XII. p. 444.*

Mas.—Lætè viridis: oculi ocellique rufo-fusci: trophi flavi; maxillæ basi et mentum viridi-ænea: antennæ fulvæ, corporis dimidio longiores, vix pubescentes; articulus 1^{us}. lætè flavus, apice fulvus; 9^{us}. et 10^{us}. fusci; clava nigra: abdomen cupreum, basi apiceque viride nitentius, subtus æneo-cupreum pubescentius: sexualia fusca: pedes lætè flavi; coxæ virides; profemora basi extus ferruginea; mesofemora intus ferrugineo vittata, spina fusca armata; protibiæ extus ferrugineo et mesotibiæ intus fusco vittatæ, hæ rubro marginatæ puncto nigro pilis nigris ciliato terminatæ; tarsi pallidè fulvi, apice fusci: alæ hyalinæ; squamulæ flavæ; nervi fulvi, ubi costam attingunt pallidè fusci; stigma concolor, parvum.

Fem.—Caput viride: antennæ pallidè fuscæ, corporis dimidio vix longiores, apice obscuriores, basi fulvæ: thorax æneo-viridis, cupreo variegatus: mesothoracis scutellum purpureo-cupreum: metathorax viridis: abdomen æneo-cupreum; segmenta basi viridia: pedes fusci; coxa æneo-virides; femora et tibiæ apice tarsique pallidè flava, hi apice fusci; protibiæ et protarsi flava, illæ extus fulvæ: alæ subhyalinæ. (Corp. long. $\frac{1}{2}$ —1 $\frac{1}{3}$ lin.; alar. $\frac{5}{4}$ —2 lin.)

Var. β.—*Mas*, caput supra et circum oculos æneo-viride: mesothoracis dorsum cupreo-viride; scutum anticè viride: metathorax æneo-viridis: abdomen apice basique æneo-viride.

Var. γ.—*Mas*, *Var. β.* similis: thorax viridis; mesothoracis scutum anticè cupreo-æneum.

Var. δ.—*Mas*, abdomen purpureo-cupreum, basi apiceque æneo-cupreum.

Var. ε.—*Mas*, caput et thorax cyaneo-viridia.

Var. ζ.—*Mas*, abdomen purpureo-cupreum, basi æneo-viride.

Var. η.—*Mas*, antennæ articulo 1^o. omninò flavo.

Var. θ.—*Mas*, thorax anticè et utrinque viridicyaneus.

Var. ι.—*Mas*, caput supra, mesothoracis scutum anticè, epimera et paraptera cupreo-ænea: abdomen cupreum, basi æneo-viride.

Var. κ.—*Mas*, caput anticè cyaneo-viride: abdomen purpureo-cupreum, basi apiceque viride.

Var. λ.—*Mas*, caput supra æneo-viride.

Var. μ.—*Mas*, prothorax omninò et mesothorax anticè æneo-virides.

Var. ν.—*Mas*, antennæ articulo 9^o. fulvo: abdomen æneo-viride.

- Var. ξ.*—*Mas*, antennæ articulis 1°. ad 7^{um}. flavis, 8°. et 9°. fulvis, 10°. fusco.
- Var. ο.*—*Mas*, nervi omnes pallidè fulvi.
- Var. π.*—*Fem.* caput supra æneo-viride : protibiæ fuscæ.
- Var. ρ.*—*Fem.* antennæ obscurè fuscæ, basi pallidiores : thorax æneo-viridis ; mesothoracis scutum cupreo-æneum.
- Var. σ.*—*Fem.* mesothoracis scutellum cupreo-æneum.
- Var. τ.*—*Fem.* metathorax æneo-viridis.
- Var. υ.*—*Fem.* femora fusco-viridia.
- Var. φ.*—*Fem.* mesothoracis scuti dorsum cyaneo-viridi maculatum.
- Var. χ.*—*Fem.* thorax æneo-viridis ; mesothoracis scutellum ferè omne et scutum posticè cupreo-cyanea.
- Var. ψ.*—*Fem.* *Var. χ.* similis sed mesothoracis macula cuprea.
- Var. ω.*—*Fem.* abdominis segmentum 1^{um}. basi viride ; 2^{um}. et sequentia basi ænea : alæ subflavescentes ; stigma obscurè fuscum.
- Var. αα.*—*Fem.* thorax cupreo-æneus : abdominis segmenta basi viridi-ænea.
- Var. ββ.*—*Fem.* thorax æneo-viridis ; metathorax viridis.

May and June ; on grass in woods ; near London. June ; Windsor. New Forest, Hampshire. Isle of Wight.

GENUS XI.—MESOPOLOBUS, *Westwood*.

Mas.—Corpus angustum, sublineare : caput parvum, thoracis latitudine : antennæ subclavatæ ; articulus 1^{us}. gracilis, sublinearis ; 3^{us}. et 4^{us}. minimi ; 5^{us}. et sequentes ad 10^{um}. subæquales, latitudine gradatim crescentes ; clava ovata, articulo 10°. paullò latior : mandibula una 4-dentata, subquadrata, vix arcuata, intus emarginata ; dens externus mediocris, 2^{us}. minor, ambo acuminati, 3^{us}. et 4^{us}. brevissimi, ille subacuminatus, hic latior obtusus ; altera similis sed dentibus 3°. et 4°. vix discretis aut conspicuis : maxillæ basi latæ, ad apices angustæ et acuminatæ ; quæque in lobum lacinia intus producta ; palpi 3-articulati ?, graciles, longitudine mediocri, articulus 1^{us}. cyathiformis, mediocris ; 2^{us}. 1½. longitudine, basi crassior ; 3^{us}. subfusiformis, 1°. et 2°. longior, ramulum intus prope basim longum gracilem emittens : labium angustum, basi conicum ; ligula minuta, anticè quasi fissa ; palpi 3-articulati, breves, crassi ; articulus 2^{us}. brevissimus : thorax elongato-ovatus, angustus : prothorax brevissimus : mesothoracis scutum et scutellum maxima ; parapsidum suturæ vix conspicuæ ; paraptera et epimera magna : metathorax mediocris : abdomen sublineare, thorace paullò brevius et angustius, basi et apice

angustatum; segmentum 1^{um}. maximum; sequentia breviora; apicalia parva: pedes graciles; metacoxæ longiores; mesotibiæ apices versus in lobum extus productæ trigonum: alæ mediocres; nervus humeralis ulnari multò longior, ramulum rejiciens nullum; cubitalis radiali brevior; stigma ramulum brevem emittens.

Sp. 1. Mesop. fasciiventris. Mas. *Viridis, antennis pedibusque flavis, abdomine cupreo flavo fasciato, alis hyalinis.*

Mesopolobus fasciiventris. *Westwood, Lond. and Edinb. Phil. Mag. Third Series. Vol. II. No. XII. p. 443.*

Lætè viridis, nitens: oculi ocellique rufo-fusci: trophi lætè flavi: mandibulæ apice rufæ: antennæ lætè flavæ, corporis dimidii longitudine: metathorax subtilissimè squameus: abdomen cupreum, ante medium latè flavo fasciatum, apice viride: sexualia fusca: pedes lætè flavi, graciles; coxæ virides; mesotibiarum lobi apice nigri et pilis nigris vestiti; tarsi apice pallidè fusci: alæ hyalinæ, subangustæ, inter nervos cubitalem et radialem fuscæ; squamulæ flavæ; nervus humeralis flavus, pilis nonnullis nigris vestitus, ubi costam attingit obscurè fuscus; ulnaris pallidè fulvus; radialis obscurior; cubitalis obscurè fuscus; stigma concolor, parvum; metalarum nervi pallidè flavi, ubi costam attingunt fulvi. (Corp. long. $\frac{2}{3}$ — $1\frac{1}{4}$ lin.; alar. $\frac{3}{4}$ — $1\frac{1}{3}$ lin.)

Var. β. — Caput, thorax et coxæ æneo-viridia: abdomen basi cupreo-æneum.

Var. γ.—*Var. β.* similis: antennæ articulis 1°. apice et 2°. supra pallidè fulvis: alæ inter nervos cubitalem et radialem subhyalinæ; nervi pallidiores.

Var. δ.—Metalarum nervi ubi costam attingunt fusci.

Var. ε.—Abdomen basi æneo-viride.

Var. ζ.—Thorax anticè et caput cyaneo-viridia: abdomen basi æneo-viride.

Var. η.—Nervus cubitalis et stigma pallidè fusca.

Var. θ.—Nervi pallidè fulvi; nervus humeralis ubi costam attingit fuscus.

May and June; on grass beneath trees; near London. June; New Forest, Hampshire. September; Isle of Wight.

GENUS XII.—EUTELUS, ^c Walker.

Mas et Fem. — Corpus multiforme : caput mediocre : antennæ clavatae ; articulus 1^{us}. *mas* subfusiformis, *fem.* sublinearis gracilior ; 3^{us}. et 4^{us}. minimi ; 6^{us}. 5^o. longior ; sequentes ad 10^{um}. gradatim breviores et latiores ; clava articulo 10^o. latior et plus duplò longior : mandibulæ 4-dentatae, subquadratae, vix arcuatae, intus emarginatae ; una dente externo mediocri acuminato, 2^o. obtusiore minore, 3^o. et 4^o. minutis, hoc latiore obtusiore ; altera dentibus brevioribus, 3^o. et 4^o. minimis vix discretis aut conspicuis : maxillæ elongatae, subarcuatae, basi latae ; intus lacinia quæque in lobum producta ; palpi 4-articulati, graciles, filiformes, longitudine mediocri ; articuli 1^{us}., 2^{us}. et 3^{us}. mediocres subæquales ; 4^{us}. subfusiformis, 2ⁱ. et 3ⁱ. longitudine : labium elongato-ovatum, posticè conicum ; ligula parva, anticè quasi fissà ; palpi 3-articulati, breves, crassi ; articulus 2^{us}. minimus : prothorax et metathorax brevia : mesothoracis scutum et scutellum maxima ; parapsidum suturæ vix conspicuæ ; paraptera et epimera benè determinata : *fem.* abdomen acuminatum ; segmentum 1^{um}. magnum ; sequentia breviora, subæqualia : oviductus plerumque occultus : pedes simplices ; mesofemora sæpè apices versus spina armata : alæ mediocres ; nervus humeralis ulnari multò longior, ramulum rejiciens nullum ; cubitalis radiali brevior ; stigma ramulum brevissimum emittens.

Sp. 1. Eut. dilectus. Mas. *Viridis, antennis pedibusque flavis, illarum articulo 8^o. clavaque fuscis, abdomine cupreo flavo fasciato, alis hyalinis.*

Lætè viridis, latus : caput thorace paullò latius : oculi ocellique fusciorum : antennæ lætè flavæ, corporis dimidii longitudine ; articulus 1^{us}. validus ; 8^{us}. pallidè fuscus ; clava fusca, apice pallidior : thorax elongato-ovatus : abdomen cupreum, ovatum, thoracis longitudine, ante medium flavo latè fasciatum, basi apiceque viride ; segmenta basi æneo-viridia : sexualia fusca : pedes lætè flavi ; coxæ virides ; tarsi apice fusci : alæ hyalinæ ; squamulæ et nervi flava, hi pilis nigris vestiti ; pro- et metalarum nervi humerales ubi costam attingunt pallidè fusci ; stigma concolor, parvum. (Corp. long. $\frac{2}{3}$ — $1\frac{1}{4}$ lin. ; alar. 1—2 lin.)

Var. β.—Antennæ articulo 2^o. basi fusco : alæ subflavescentes.

Var. γ.—Var. β. similis: alæ hyalinæ; stigma flavum.

Var. δ.—Abdominis segmenta basi cuprea: stigma flavum.

Var. ε.—Metathorax basi æneus.

Var. ζ.—Caput et thorax æneo-viridia: abdomen purpureo-cupreum: stigma flavum.

June to October; on grass beneath trees; near London.
June; Windsor. September; Linton, North Devonshire.
New Lanark, Scotland.

Sp. 2. *Eut. immaculatus.* Mas. *Viridis, antennis pedibusque flavis, illarum articulo 8º. clavaque fuscis, abdomine cupreo, alis hyalinis.*

Lætè viridis, crassus, *E. dilecti* statura: caput thorace paullò latius: oculi ocellique rufo-fusci: antennæ lætè flavæ, corporis dimidii longitudine; articulus 1^{us}. validus; 8^{us}. pallidè fuscus; clava fusca, apice pallidior: abdomen cupreum, immaculatum, thorace paullò brevius et angustius, basi apiceque cupro-æneum: sexualia fusca: pedes læte flavi; coxæ virides; tarsi apice pallidè fusci; protibiæ apice et protarsi fulvo-flava: alæ hyalinæ; squamulæ flavæ; nervi pallidè fulvi; stigma minutum. (Corp. long. 1 lin.; alar. 1½ lin.)

Var. β.—Metathorax æneo-viridis.

August; on grass beneath trees; near London.

Sp. 3. *Eut. signatus.* Mas. *Viridis, E. dilecto angustior, antennis pedibusque flavis, illarum articulo 8º. clavaque fuscis, abdomine cupreo flavo fasciato, alis hyalinis.*

Viridis, *E. dilecto* minor angustior et multò obscurior: caput thorace latius: oculi ocellique rufo-fusci: antennæ pallidè flavæ, corporis dimidii longitudine; articuli 2^{us}. supra, 7^{us}. apice et 8^{us}. fusci; clava nigro-fusca, apice basique flava: abdomen cupreum, thorace paullò brevius, flavo antè medium latè fasciatum: sexualia fusca: pedes pallidè flavi; coxæ virides; tarsi apice pallidè fusci: alæ hyalinæ; squamulæ flavæ; nervi pallidè fulvi; stigma parvum. (Corp. long. ½ lin.; alar. ⅔ lin.)

September; Ambleside, Westmoreland.

Sp. 4. Eut. pygmæus. Mas. *Viridis, E. dilecto angustior, antennis fulvis, articulo 2º. clavaque fuscis, abdominis fascia pedibusque flavis, alis hyalinis.*

Viridis, *E. dilecto* angustior et minor: caput thoracis latitudine: antennæ fulvæ, corporis dimidio paullò longiores; articulus 1^{us}. lætè flavus; 2^{us}. fuscus, apice flavescens; clava fusca; oculi ocellique rufo-fusci: abdomen thoracis longitudine, ante medium flavo fasciatum: pedes lætè flavi; coxæ virides; tarsi apice pallidè fusci; alæ hyalinæ; squamulæ et nervi pallidè flava; stigma minutum. (Corp. long. $\frac{1}{3}$ lin.; alar. $\frac{1}{2}$ lin.)

August; on grass beneath trees; near London.

Sp. 5. Eut. diffinis. Mas. *Viridis, antennis fulvis apice fuscis, abdomine fulvo fasciato, pedibus flavis, alis hyalinis, metalis fusco maculatis.*

Lætè viridis, *E. dilecto* paullò angustior: caput thorace vix latius: oculi ocellique rufo-fusci: antennæ fulvæ, corporis dimidii longitudine; articulus 1^{us}. flavus; 2^{us}. basi, 3ⁱ. et sequentium ad 8^{um}. suturæ et clava fusca, hæc apice pallidior: abdomen æneo-viride, thorace paullò brevius, basi apiceque viride; segmentum 2^{um}. apice fulvum: pedes lætè flavi; coxæ virides; mesotibiæ intus et tarsi apice fusca: alæ hyalinæ; squamulæ et nervi flava; stigma minutum: metalæ nisi ad basim subfusca; maculæ in cujusque disco fusca irregulares medio connectæ duæ. (Corp. long. $\frac{1}{3}$ — $\frac{3}{4}$ lin.; $\frac{1}{2}$ —1 lin.)

Var. β.—Caput et thorax æneo-viridia.

Var. γ.—Mesotibiæ intus ad apices obscurè fusca, ante apices flavæ.

Var. δ.—Proalæ sub nervum ulnarem fulvescentes.

August to October; on grass in fields; near London. September; Isle of Wight; Lyme Regis, Dorsetshire; Sidmouth and Plymouth, Devonshire. New Lanark, Scotland.

Sp. 6. Eut. jucundus. Mas. *Viridis, antennis pedibusque flavis, abdomine cupreo flavo fasciato, alis hyalinis.*

Lætè viridis, angustus: caput thorace vix latius: oculi ocellique rufo-fusci: antennæ lætè flavæ, corporis dimidio paullò breviores: caput posticè thoracisque latera et apex æneo-viridi variegata: abdomen cupreum, thorace paullò brevius, ante medium latè

flavo-fasciatum, apice viride: pedes lætè flavi; coxæ æneo-virides; tarsi apice pallidè fusi: alæ hyalinæ; squamulæ pallidè flavæ; nervi pallidè fulvi; stigma minutum. (Corp. long. $\frac{3}{4}$ — $1\frac{1}{4}$ lin.; alar. $1\frac{1}{4}$ — $1\frac{3}{4}$ lin.)

Var. β.—Caput et thorax omnino viridia.

Var. γ.—Abdomen basi æneo-viride.

May and June; on grass beneath trees; near London.

Sp. 7. Eut. placidus. Mas. *Viridis*, E. diffinis *statura*, *antennis flavis fusco maculatis*, *pedibus flavis*, *alis hyalinis*.

Lætè viridis: caput thorace vix latius: oculi ocellique rufo-fusci: antennæ fuscae, corporis dimidio paullò longiores; articuli 1^{us}., 3^{us}., 4^{us}., 5^{us}., et 8^{us}. flavi; 9^{us}. et 10^{us}. pallidè fusi: abdomen thorace paullò longius; discus cupreo-viridis: sexualia fusca: pedes lætè flavi; coxæ virides; tarsi apice pallidè fusi; protarsi fulvi: alæ hyalinæ; squamulæ et nervi pallidè flava; stigma minutum. (Corp. long. $\frac{3}{4}$ lin.; alar. 1 lin.)

Var. β.—Antennæ articulo 8^o. supra pallidè fusco.

Var. γ.—Mesothoracis paraptera et epimera æneo-viridia: antennæ articulo 8^o. pallidè fusco: abdominis discus obscurè cupreus.

Var. δ.—Antennæ articulis 6^o. et 7^o. pallidè fuscis; 8^o., 9^o. et 10^o. fulvis: thoracis latera abdominisque margo æneo-viridia.

August; on lime-trees; near London.

Sp. 8. Eut. ocellus. Mas. *Viridi-æneus*, *antennis nigris*, *abdominis fascia flava*, *pedibus fulvis*, *pro-alis quasi ocelligeris*.

Viridi-æneus, obscurus, latus: caput thorace paullò latius: oculi ocellique rufo-fusci: trophi fusi: antennæ nigræ, corporis dimidii longitudine, apice nigro-fuscae; articulus 1^{us}. pallidè flavus; 2^{us}., 3^{us}. et 4^{us}. pallidè fusi: abdomen cupreum, thoracis longitudine, ante medium flavo-fasciatum, apice viridescens: pedes fulvi, validi; coxæ cyaneo-virides; genua, tibiæ basi, meso- et metatarsi flava, hi apice pallidè fusi: alæ subfulvescentes, angustæ, breves, infectæ?; proalæ fusco apice marginatæ et sub stigmatè maculatæ; squamulæ et nervi pallidè fusca, nervus ulnaris obscurior; stigma obscurè fuscum, mediocre. (Corp. long. $1\frac{1}{4}$ lin.; alar. $1\frac{1}{2}$ lin.)

New Lanark, Scotland.

Sp. 9. Eut. eximius. Fem. *Æneus aut viridis, antennis pedibusque fulvis, alis hyalinis plus minusve flavescens*.

Lætè æneus, latus: caput thoracis latitudine, posticè viride: oculi ocellique rufo-fusci: antennæ fulvæ, subtus pallidiores, corporis dimidio vix breviores; articulus 1^{us}. pallidè fulvus; sequentes basi fusci: thorax ovatus; segmentorum margines virides: abdomen cupreo-æneum, thorace paullò brevius, subtus non angulatum; discus obscurè cupreus: pedes fulvi; coxæ viridi-æneæ; genua, tibiæ apice tarsique flava; hi apice fusci; protibiæ et protarsi fulvo-flava: alæ hyalinæ, sub costam flavescens; squamulæ et nervi flava; stigma minutum. (Corp. long. 1—1½ lin.; alar. 1¾—2¼ lin.)

Var. β.—Æneo-viridis: abdomen basi viride; segmenta 1^o. ad 4^{um}. purpureo-cuprea.

Var. γ.—Lætè viridis: abdomen æneum, basi lætè viride, medio cupreum: coxæ virides; profemora et protibiæ flava: alæ vix flavescens.

September and October; on oak-trees; near London. New Lanark, Scotland.

Sp. 10. Eut. platycerus. Fem. *Æneus, antennis pedibusque fuscis, tarsis pallidè flavis, alis hyalinis plus minusve flavescens*.

Æneus, latus, parùm nitens: caput æneo-viride, thoracis latitudine: oculi ocellique rufo-fusci: antennæ fuscae, validæ, corporis dimidii longitudine, subtus pallidiores; articulus 1^{us}. omninò 2^{us}.-que subtus fulvi: abdomen obscurè cupreum, thoracis longitudine, basi lætè æneo-viride, subtus non angulatum; segmenta basi viridia: pedes pallidè fusci; coxæ æneo-virides; trochanteres, femora apice, tibiæ apice et basi tarsique pallidè flava; hi apice fusci; protibiæ et protarsi fulva: alæ hyalinæ, subflavescens; squamulæ fulvæ; nervi flavi; stigma fulvum, parvum. (Corp. long. 1—1¼ lin.; alar. 1½—1¾ lin.)

Var. β.—Caput viride: thorax viridi-æneus.

Var. γ.—Caput viridi-æneum: thorax cupreo-æneus; segmentorum margines virides.

Var. δ.—Caput viride: thorax cupreus: abdominis segmenta basi ænea.

Var. ε.—Thorax viridi-æneus, cupreo variegatus.

Var. ζ.—Thorax æneo-cupreus; segmentorum margines virides: abdominis segmenta basi viridi-ænea.

Var. η.—Caput viride: thorax viridi-æneus; mesothoracis scutellum cupreo-æneum.

Var. θ.—Caput cyaneo-viride: thorax æneo-cupreus: meso- et metapedum femora et tibiæ fusca: alarum nervi fulvi; stigma fuscum.

Var. ι.—Metathorax æneo-viridis.

Var. κ.—Abdominis segmenta basi ænea.

Var. λ.—Antennæ articulo 1^o. pallidè fusco: alarum nervi fulvi; stigma fuscum.

Var. μ.—Tibiæ omnes fulvæ.

Var. ν.—Protibiæ et protarsi flava.

June to October; on grass beneath trees; near London.
June and September; Isle of Wight. June; New Forest, Hampshire. September; Sidmouth, Devonshire.

Sp. 11. Eut. bicolor. Fem. *Præcedentis ferè statura, plerumque major, colore lætiore, pedibus pallidioribus.*

Æneus: caput viride, thorace sublatius: oculi ocellique rufo-fusci: antennæ pallidè fuscæ, corporis dimidii vix longitudine; articulus 1^{us}. omninò 2^{us}.-que subtus fulvi: mesothoracis scutellum cupreo-æneum: abdomen æneum, thorace paullò longius; segmentum 1^{um}. basi æneo-viride cupreo maculatum; sequentia apice obscurè cuprea: pedes fulvi; coxæ viridi-æneæ; genua, tibiæ apice et tarsi flava; hi apice obscurè fusci; protibiæ et protarsi pallidè fulva; alæ hyalinæ, minimè flavescentes; squamulæ et nervi fulva; stigma paullò obscurius, parvum. (Corp. long. $1\frac{1}{4}$ — $1\frac{1}{2}$ lin.; alar, $1\frac{3}{4}$ —2 lin.)

Var. β.—Abdominis segmenta basi cupreo-ænea.

Var. γ.—Femora, meso- et metatibiæ pallidè fusca.

September; on grass in fields; near London. Isle of Wight.

Sp. 12. Eut. platynotus. Fem. *Viridis, E. platyceri statura.*

Viridis: caput cyaneo-viride, thorace vix latius: oculi ocellique rufo-fusci: antennæ fulvæ, corporis dimidii longitudine, supra pallidè fuscæ, apice obscuriores; articulus 1^{us}. fulvus: abdomen thoracis longitudine; segmenta apice cupreo-ænea: pedes fusci; coxæ virides; tibiæ apice tarsique straminea, hi apice fusci;

trochanteres, genua, protibiæ et protarsi fulva: alæ hyalinæ; squamulæ et nervi fulva; stigma parvum. (Corp. long. $\frac{3}{4}$ —1 lin.; alar. $1\frac{1}{4}$ — $1\frac{1}{2}$ lin.)

Var. β.—Thorax posticè æneo-viridis: protibiæ et protarsi flava.

July; on grass beneath trees; near London.

Sp. 13. Eut. sobrinus. Fem. Præcedentibus 3 paullo longior.

Æneus: caput viride, thorace sublatius: oculi ocellique rufo-fusci: antennæ obscurè fulvæ, corporis dimidii longitudine, supra fuscae; clava obscurior; articulus 1^{us}. fulvus, apice fuscus: thoracis latera viridi-ænea: abdomen obscurè cupreum, thorace paullo longius; segmenta basi lætè æneo-viridia: pedes fusci; coxæ æneæ; femora viridi-fusca; meso-femora apices versus spina fusca subtus armata; trochanteres et protarsi fulva; genua, tibiæ apicè tarsique flava, hi apice fusci; protibiæ flavæ, extus fuscae: alæ hyalinæ, minimè fulvescentes; squamulæ et nervi fulva; stigma obscurius, parvum. (Corp. long. 1— $1\frac{1}{4}$ lin.; alar. $1\frac{1}{2}$ — $1\frac{3}{4}$ lin.)

Var. β.—Caput inter oculos æneo-viride: antennæ articulis 1^o, 3^o. 4^o.-que omninò fulvis: thorax viridi-æneus: protibiæ extus fulvæ.

Var. γ.—Caput inter oculos æneo-viride: antennæ fuscae; articulus 1^{us}. fulvus: thoracis dorsum cupreo-æneum: abdominis segmenta 2^{um}. et sequentia basi ænea.

Var. δ.—Antennæ articulo 1^o omninò fulvo; thorax supra cupreo-æneus; segmentorum margines virides: abdominis segmentum 1^{um}. basi cupreo variegatum.

Var. ε.—Thorax supra abdominisque segmenta 2^{um}. et sequentia basi cupreo-ænea; protibiæ utrinque fuscae; pro-alæ fulvescentes.

Var. ζ.—Mesothoracis scutellum cupreo-æneum: abdominis segmenta basi viridi-ænea.

May and June; on grass in fields; near London. June; Windsor.

Sp. 14. Eut. catenatus. Fem. Viridis, 4-præcedentibus angustior.

Viridis: caput thoracis latitudine: oculi ocellique rufo-fusci: antennæ fusco-fulvæ, corporis dimidii longitudine; articulus 1^{us}. fulvus: mesothoracis scutum æneo-viride: abdomen cupreum, thorace paullo longius; segmenta basi æneo-viridia: pedes fulvi; coxæ virides; femora, meso- et metatibiæ fusca, apice flava;

meso- et metatarsi flavi, apice fusci: alæ hyalinæ, nonnunquam minimè flavescentes; squamulæ et nervi fulva; stigma parvum. (Corp. long. $\frac{5}{4}$ —1 lin.; alar. $1\frac{1}{4}$ — $1\frac{1}{2}$ lin.)

Var. β.—Caput inter oculos thoracisque dorsum æneo-viridia.

June; on grass beneath trees; near London; New Forest, Hampshire.

Sp. 15. Eut. inornatus. Fem. *Æneus, antennis nigris, pedibus fuscis, alis subfuscis.*

Obscurè æneus, brevis: caput nigrum, thorace vix latius: oculi ocellique rufo-fusci: antennæ nigræ, corporis dimidio breviores; articulus 1^{us}. basi fuscus: abdomen cupreo-æneum, thorace latius sed non longius, subtus non angulatum; segmentum 1^{um}. basi et sequentia apice viridi-ænea: pedes fusci; coxæ æneæ; femora apice, tibiæ basi, trochanteres et tarsi fulva, hi apice fusci: alæ subfuscae; squamulæ et nervi fusca; stigma minutum. (Corp. long. $\frac{5}{4}$ lin.; alar. $1\frac{1}{6}$ lin.)

Var. β.—Caput, thorax utrinque et posticè, coxæ et femora obscurè viridia.

Var. γ.—Abdomen basi cupreo-viride.

September; Westmoreland and Cumberland.

Sp. 16. Eut. fulvicornis. Fem. *Æneo-viridis, E. dilecto angustior, antennis fulvis, pedibus flavis, alis hyalinis.*

Lætè æneo-viridis: caput thoracis latitudine: oculi ocellique rufo-fusci: antennæ fulvæ, corporis dimidio breviores; clava pallidior; articulus 1^{us}. flavus: abdomen thorace longius, subtus paullò angulatum, apice acuminatum vix attenuatum; discus et apex obscurè cuprei: oviductus fulvus: pedes lætè flavi; coxæ virides; tarsi apice fusci; protarsi fulvo-fusci: alæ hyalinæ, sub costam minimè flavescents; squamulæ et nervi flava; horum humeralis ubi costam attingit et cubitalis fulvi; stigma minutum. (Corp. long. 1— $1\frac{1}{2}$ lin.; alar. $1\frac{1}{2}$ — $2\frac{1}{2}$ lin.)

Var. β.—Caput et thorax viridia, hujus segmentorum margines æneo-virides.

Var. γ.—Caput et thorax omninò viridia.

Var. δ.—Cyaneo-viridis: abdomen viride; discus cupreus.

Var. ε.—Antennæ articulo 1^o. apice supra pallidè fulvo.

Var. ζ.—Abdomen cupreo-æneum; discus obscurè purpureo-cupreus.

Var. η.—Abdomen basi cupreo-viride.

Var. θ.—Abdomen supra obscurè cupreum, basi cupreo-viride, subtus et utrinque æneum segmentis basi cupreis.

Var. ι.—Nervi omninò flavi.

Var. κ.—*Var. ι* similis : metafemora fulva.

Var. λ.—*Var. κ* similis : abdomen cupreum, basi lætè viride ; segmenta utrinque et subtus basi viridia.

Var. μ.—*Var. κ* similis : antennæ fusco-fulvæ.

Var. ν.—Nervi radialis et ulnaris fulvi.

July to October ; on grass in fields ; near London. September ; Isle of Wight ; Linton, Devonshire ; New Lanark, Scotland.

Sp. 17. Eut. flavipes. Fem. *Viridis, antennis fuscis, pedibus flavis, alis hyalinis.*

Lætè viridis : caput thorace paullò latius : oculi ocellique rufo-fusci : antennæ fuscae, corporis dimidii longitudine ; articulus 1^{us}. basi fulvus ; 2^{us}. supra obscurè fuscus : thorax brevis : abdomen longius et gracilius, basi angustum, medio supra obscurè cupreum, subtus æneum non angulatum : pedes flavi ; coxæ virides ; tarsi apice fusci ; protarsi fulvi : alæ hyalinæ ; squamulæ et nervi flava ; stigma fulvum, minutum. (Corp. long. $\frac{3}{4}$ lin. ; alar. $1\frac{1}{4}$ lin.)

July ; on grass in fields ; near London.

Sp. 18. Eut. æquus. Fem. *Æneo-viridis, antennis fusco-fulvis, pedibus flavis, alis hyalinis.*

Æneo-viridis, angustus, sublinearis : caput thorace latius : oculi ocellique rufo-fusci : antennæ fusco-fulvæ, corporis dimidio breviores ; articulus 1^{us}. pallidè fulvus : abdomen æneum, thorace paullò brevius et latius, basi cupreo-viride, subtus non angulatum, apice vix acuminatum : pedes lætè flavi ; coxæ æneo-virides ; trochanteres pallidè fulvi ; meso- et metatarsi straminei, apice fusci : alæ hyalinæ, minimè flavescentes ; squamulæ et nervi flava ; stigma minutum. (Corp. long. $1\frac{1}{4}$ — $1\frac{1}{2}$ lin. ; alar. $1\frac{1}{2}$ —2 lin.)

Var. β.—Viridis : antennæ articulo 1^o. supra apice fusco-fulvo : abdominis discus cupreus.

Var. γ.—Antennæ fuscae, apice fusco-fulvæ : abdomen æneo-viride.

Var. δ.—Antennæ fuscae ; articulus 1^{us}. fulvus, basi flavus : abdomen viride, basi nitentius ; discus cupreus : metathorax lætè viridis.

Var. ε.—Abdomen cupreo-æneum ; discus obscurè cupreus.

Var. ζ.—Thoracis dorsum cupreo et purpureo maculatum.

Var. η.—Femora omnia et mesofemora præcipuè basi fulva.

August to October; on grass in fields; near London. September; Isle of Wight. New Lanark, Scotland.

Sp. 19. Eut. planus. Fem. *Cyaneo-viridis, antennis nigris, pedibus flavis, alis hyalinis.*

Lætè cyaneo-viridis, angustus, elongatus: caput viride, thorace vix latius: oculi ocellique rufo-fusci: antennæ nigræ, corporis dimidio breviores, apice fuscæ; articulus 1^{us}. flavus, apice nigro-fuscus: abdomen æneum, thorace paullò brevius, subtus non angulatum; discus cupreo-æneus: pedes lætè flavi; coxæ virides; meso- et metapedum tibiæ tarsique straminea, hi apice pallidè fusci: alæ hyalinæ, apice subgriseæ; squamulæ et nervi flava; stigma minutum. (Corp. lon. $1\frac{1}{2}$ lin.; alar. $2\frac{1}{4}$ lin.)

May; on grass beneath trees; near London.

Sp. 20. Eut. gracilis. Fem. *Viridis, antennis nigris, pedibus flavis, alis hyalinis.*

Lætè viridis, angustus, elongatus: caput thorace vix latius; oculi ocellique rufo-fusci: antennæ nigræ, corporis dimidio breviores, apice nigro-fuscæ; articulus 1^{us}. fulvus, apice nigro-fuscus: mesothoracis scutellum æneo-viride: abdomen thorace paullò longius, subtus non angulatum; discus cupreus; segmenta basi subtus et utrinque ænea: pedes lætè flavi; coxæ virides; tarsi apice fusci: alæ hyalinæ; squamulæ flavæ; nervi pallidè fulvi; stigma minutum. (Corp. long. $1\frac{1}{2}$ lin.; alar. 2 lin.)

New Lanark, Scotland.

Sp. 21. Eut. helvipes. Fem. *Viridis, pedibus flavis, antennis femoribusque fulvis, alis hyalinis.*

Lætè viridis, longus, angustus: caput thoracis latitudine: oculi ocellique fusco-rufi: trophi flavi: antennæ fulvæ, corporis dimidio breviores, supra fuscæ; articulus 1^{us}. omninò fulvus: abdomen cupreum, angustum, thorace longius, subtus non angulatum; segmentum 1^{um}. basi lætè viride; sequentia basi æneo-viridia: pedes lætè flavi; coxæ virides; trochanteres pallidè fulvi; femora fulva; metafemora fusca; tarsi apice fusci: alæ hyalinæ; squamulæ et nervi pallidè flava; stigma minimum. (Corp. long. $1\frac{1}{4}$ lin.; alar. $1\frac{3}{4}$ lin.)

New Lanark, Scotland.

Sp. 22. Eut. posticus. Fem. *Æneo-cupreus, antennis nigris, pedibus fulvo-flavis, alis hyalinis.*

Æneo-cupreus, obscurus: caput viridi-æneum, thoracis latitudine: oculi ocellique rufo-fusci: antennæ nigræ, corporis dimidio paullò breviores, apice nigro-fuscæ; articulus 1^{us}. basi fulvus: thorax angustus; metathorax lætè viridis: abdomen purpureo-cupreum, thorace latius sed non longius, apice et subtus viridi-æneum; segmentum 1^{um}. cupreo-viride; reliqua basi viridia: pedes flavi; coxæ virides; femora basi fusca; profemora basi fulva; meso- et metatibiæ et protarsi fulva, illæ basi apiceque flavæ; tarsi apice fusci: alæ minimè fuscæ, parvæ; squamulæ et nervi fulva; stigma fuscum, parvum. (Corp. long. $1\frac{1}{4}$ lin.; alar. $1\frac{2}{3}$ lin.)

New Lanark, Scotland.

Sp. 23. Eut. elevatus. Fem. *Æneo-viridis, antennis nigris, pedibus fulvis, femoribus viridibus, alis subhyalinis.*

Æneo-viridis, obscurus: caput thoracis vix latitudine: oculi ocellique rufo-fusci: antennæ nigræ, corporis dimidio paullò breviores, apice nigro-fuscæ; articulus 1^{us}. viridis, basi fulvus; 2^{us}. æneo-viridis: abdomen thorace longius, subtus angulatum, apice elevatum acuminatum attenuatum; segmenta apice viridia: pedes fulvi; coxæ æneo-virides; femora viridia, apice fulva; trochanteres et protarsi fusci; meso- et metapedum tibiæ apice tarsique basi flava: alæ subhyalinæ; squamulæ et nervi fulva; stigma obscurius, parvum. (Corp. long. $1\frac{1}{4}$ —2 lin.; alar. $1\frac{2}{3}$ — $2\frac{2}{3}$ lin.)

Var. β.—Meso- et metafemora fusco cingulata.

Var. γ.—Viridi-æneus: abdomen basi æneum.

Var. δ.—Antennæ nigro-fuscæ: abdomen æneo-viride; segmenta basi cuprea.

September; Isle of Wight.

Sp. 24. Eut. intermedius Fem. *Præcedenti similis; abdomen paullò brevius et latius.*

Viridi-æneus, obscurus: caput æneo-viride, thoracis latitudine: oculi ocellique rufo-fusci: antennæ fuscæ, corporis dimidii longitudine; articulus 1^{us}. basi fulvus: abdomen lætè viride, thorace vix longius, apice basique cupreo variegatum, medio cupreum: pedes flavi; coxæ et femora æneo-viridia; tibiæ fusco fasciatae; tarsi apice fusci; trochanteres, protibiæ et protarsi fulva: alæ subhy-

alinæ; squamulæ et nervi fulva, horum humeralis flavus; stigma parvum, obscurius. (Corp. long. $1\frac{1}{2}$ lin.; alar. 2 lin.)

Var. β.—Meso- et metatibiæ fulvo fasciatæ: alarum squamulæ fulvæ; nervi flavi; stigma fuscum.

September; Isle of Wight.

Sp. 25. Eut. semotus. Fem. *Viridi-æneus, antennis fuscis, pedibus flavis fusco cingulatis, alis hyalinis.*

Viridi-æneus, obscurus, *E. clavato* et *intermedio* abdomine minus attenuato hoc quoque antennis gracilioribus discretus: caput æneo-viride, thoracis latitudine: oculi ocellique rufo-fusci: antennæ obscurè fuscæ, corporis dimidii longitudine; articulus 1^{us}. fusco-viridis, basi fulvus: abdomen purpureo-cupreum, apice elevatum cupreo-viride, subtus angulatum cupreum; segmentum 1^{um}. cupreum, basi viride; 2^{um}. et sequentia ad 5^{um}. apice ænea: pedes flavi; coxæ virides; trochanteres fusci; femora fusco-viridia; tibiæ fusco cingulatæ; tarsi apice fusci; protibiæ et protarsi fulva: alæ hyalinæ; squamulæ fulvæ; nervi flavi; stigma fuscum, parvum. (Corp. long.; $1\frac{1}{2}$ — $1\frac{2}{3}$ lin.; alar. 2— $2\frac{1}{4}$ lin.)

Var. β.—Antennæ articulo 1^o. fusco basi fulvo: abdominis segmentum 1^{um}. basi cupreo maculatum, 5^{um}. basi æneo-viride: meso- et metatibiæ fulvo cingulatæ.

September; Isle of Wight.

Sp. 26. Eut. altus. Mas et Fem. *Viridis, antennis fuscis, pedibus fulvo flavis, femoribus viridibus, alis subhyalinis.*

Mas.—Cyaneo-viridis, crassus: caput thorace latius: oculi ocellique rufo-fusci: antennæ fuscæ, corporis dimidii longitudine; articulus 1^{us}. fulvus, apice fuscus: abdomen thoracis longitudine; discus cupreus: sexualia fusca: pedes lætè flavi; coxæ virides; femora viridi-fusca, apice basique flava; tarsi apice fusci; protarsi fulvi: alæ subhyalinæ; squamulæ et nervi pallidè fusca; stigma parvum.

Fem.—Lætè viridis: antennæ corporis dimidio breviores: abdomen cyaneo-viride; segmenta basi cuprea: trochanteres fusci; femora viridia, apice basique fusca; tibiæ fulvæ, apice flavæ; metatibiæ fusco cingulatæ; protarsi fusci. (Corp. long. 1— $1\frac{1}{4}$ lin.; alar. $1\frac{1}{2}$ — $1\frac{3}{4}$ lin.)

Var. β.—*Fem.* antennæ obscurè fuscæ; articulus 1^{us}. basi fulvus: abdominis discus cupreus: tibiæ fuscæ, apice basique fulvæ; squamulæ et nervi obscurè fusca.

June; Windsor Forest.

Sp. 27. Eut. chlorospilus. Mas et Fem. *Viridis aut æneus, antennis nigris aut fuscis, maris abdomine flavo maculato, pedibus fuscis, femoribus nonnunquam viridibus, tarsis flavis, alis subhyalinis.*

Mas.—Æneus, obscurus: caput æneo-viride, thorace latius: oculi ocellique fusci: antennæ fuscae, corporis dimidii longitudine; articulus 1^{us}. fulvus, apice fuscus: abdomen obscurè æneo-viride, thoracis longitudine, basi viride nitentius, ante medium flavo maculatum: sexualia pallida: pedes fusci; coxæ æneo-virides; femora viridi-fusca; tibiæ apice basique flavæ; tarsi flavi, apice fusci: alæ subhyalinæ; squamulæ et nervi pallidè fusca; stigma mediocre.

Fem.—Viridi-æneus: antennæ nigro-fuscae, corporis dimidio breviores; articulus 1^{us}. niger, basi fuscus: abdomen thorace paullo longius, subtus angulatum, basi lætè viride: oviductus fulvus: pedes flavi; coxæ viridi-æneæ; femora fusca; tibiæ fusco cingulatæ; tarsi apice fusci: protibiæ et protarsi fulvo-flava. (Corp. long. $\frac{3}{4}$ —1 lin.; alar. $1\frac{1}{4}$ — $1\frac{1}{2}$ lin.)

Var. β.—*Fem.* Viridis: antennæ nigræ; articulus 1^{us}. nigro-viridis, basi fuscus: abdomen cupreum, basi, apice et subtus viride: coxæ et femora viridia; trochanteres et tibiæ fusca.

September; Isle of Wight. Exeter, Devonshire.

Sp. 28. Eut. fuscipennis. Fem. *Æneo-viridis, antennis nigro-fuscis, pedibus fuscis, femoribus viridibus, alis fuscis.*

Obscurè æneo-viridis: caput obscurè viride, thorace latius: oculi ocellique rufo-fusci: antennæ nigro-fuscae, corporis dimidio breviores, apice subtus pallidiores; articulus 1^{us}. fulvus: mesothorax posticè et metathorax virides: abdomen obscurè cupreum, thorace paullo longius, subtus angulatum, basi lætè viride, apice æneo-viride; segmenta utrinque basi viridia: pedes fusci; coxæ et femora viridia, hæ apice basique fusca; tibiæ viridi-fuscae, apice flavæ; tarsi flavi, apice fusci; protibiæ et protarsi fulva, hi apice fusci, illæ apice flavæ: alæ fuscae, basi pallidiores; squamulæ et nervi obscurè fusca; stigma mediocre. (Corp. long. $1\frac{1}{2}$ — $1\frac{2}{3}$ lin.; alar. 2— $2\frac{1}{4}$ lin.)

Var. β.—Antennæ articulo 1^o. apice fusco: thorax viridis: abdominis segmenta supra basi cyaneo-viridia.

Var. γ.—Metathorax cupreo-viridis.

June; Windsor Forest.

